



ISSN: 2617-6548

URL: [www.ijirss.com](http://www.ijirss.com)

## The UV and FTIR Fingerprint of *Ocimum kilimandscharicum* Guerke Essential oil: A Eugenol-Rich Chemo Type

 Rabia Ayoubi<sup>1</sup>,  Sadia Wali<sup>2</sup>,  Gyanesh B. Singh<sup>3\*</sup>

<sup>1,3</sup>Department of Biotechnology, School of Bioengineering and Biosciences, Lovely Professional University, Punjab, India.

<sup>2</sup>Department of Pharmacognosy, Faculty of Pharmacy, Kabul University, Kabul, Afghanistan.

\*Corresponding author: Gyanesh B. Singh (Email: [matechnology@gmail.com](mailto:matechnology@gmail.com))

### Abstract

*Ocimum kilimandscharicum* Guerke, a member of the family Lamiaceae, is a valuable medicinal plant used both in traditional and modern medicine. It is a perennial aromatic undershrub with tremendous phytochemical polymorphism. The present study aims to assess the amount of eugenol in the essential oil (EO) of *O. kilimandscharicum*. Eugenol is one of the most popular phenolic compounds, which is naturally synthesized and extracted from the EO of different plant species. The fresh leaves and flowers of *O. kilimandscharicum* were used to extract EO using a hydrodistillation method. Ultraviolet (UV) and Fourier Transform Infrared (FTIR) spectrometry techniques were used to assess and quantify the chemical fingerprint of the EO and their main phytoconstituents. In this study, eugenol showed its peak absorbance to be around 282 nm in both the EO and pure eugenol spectra. The FTIR spectra of the EO and eugenol showed different functional groups determined by comparing the vibration frequencies in wave numbers of the EO and eugenol spectra with those of an IR correlation chart. Eugenol is a well-known phenolic compound with medicinal and economic value. The UV and FTIR spectra of the EO of *O. kilimandscharicum* proved the presence of a high amount of eugenol in the *O. kilimandscharicum* plant.

**Keywords:** *Ocimum kilimandscharicum* guerke, Medicinal plants, Essential oil, Eugenol, UV spectrometry, FTIR spectrometry.

**DOI:** 10.53894/ijirss.v5i1.307

**Funding:** This study received no specific financial support.

**History: Received:** 18 November 2021/**Revised:** 15 December 2021/**Accepted:** 28 December 2021/**Published:** 6 January 2022

**Copyright:** © 2022 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Acknowledgment:** All authors contributed equally to the conception and design of the study.

**Competing Interests:** The authors declare that they have no competing interests.

**Transparency:** The authors confirm that the manuscript is an honest, accurate, and transparent account of the study was reported; that no vital features of the study have been omitted; and that any discrepancies from the study as planned have been explained.

**Ethical:** This study follows all ethical practices during writing.

**Publisher:** Innovative Research Publishing

### 1. Introduction

Phenylpropenes contain a large number of organic molecules, which plants produce for purposes such as defensive compounds against animal predators and microorganisms, structural components, growth, and floral attractants for the pollinators. In addition, these compounds have long been used by humans as food preservatives, food additives, medicinal agents, etc. Chavicol, t-anol, eugenol, isoeugenol, and methyl eugenol are well-known examples of phenylpropenes currently used for industrial and pharmaceutical purposes [1, 2].

Eugenol (4-allyl-2-methoxyphenol), with the molecular formula of C<sub>10</sub>H<sub>12</sub>O<sub>2</sub>, is an organic phenolic compound that is naturally synthesized and extracted from the essential oil of different plant species such as clove, basil, cinnamon, lemon balm, and nutmeg [3, 4]. This phytoconstituent is known for its medicinal value as an antiseptic [5], dental anesthetic [6], antioxidant [3, 7], anti-inflammatory [8], antiviral [9], antimicrobial [10], antibacterial, antifungal [11], anthelmintic [12,

13] etc. In addition, eugenol is used in the perfumery industry, food industry as a food additive [5] in clove-flavored cigarettes, dental and oral hygiene preparations, as an irritant, sensitizer, and in the production of local anesthetics [5, 14].

The genus *Ocimum* “L”. is one of the most important genera of the family Lamiaceae, which includes a large number of medicinal aromatic plants. These plants are widely distributed through the tropical and warmer temperate regions of Asia, Africa, and Central and Southern America [15-17]. Due to their medicinal and economic importance, these plants are cultivated worldwide. The aromatic herbs, under- shrubs, and shrubs of genus *Ocimum*, including *O. kilimandscharicum*, produce essential oils (EOs) of different aromas and chemical profiles with tremendous value in pharmaceutical, perfume, and food industries [15, 18-21].

*Ocimum kilimandscharicum* is a perennial aromatic undershrub native to Central and East African countries, including Rwanda, Kenya, Tanzania, Uganda, Sudan, and Ethiopia [22]. It can be cultivated in flat and hilly areas. Due to its geographic distribution and abundance in African countries, *O. kilimandscharicum* is commonly called African blue basil. The plant has attracted attention due to its high camphor content and as an exotic plant.

Taxonomical classification of *Ocimum kilimandscharicum* Guerke

Kingdom: Plantae

Subkingdom: Tracheobionta

Division: Magnoliophyta

Subdivision: Spermatophyta

Class: Magnoliopsida

Subclass: Asteridae

Order: Lamiales

Family: Lamiaceae

Genus: *Ocimum*

Species: *O. kilimandscharicum* Guerke

Synonyms: *O. johnstonii* Baker , and *O. tortuosum* Baker.



**Figure 1.**  
*Ocimum kilimandscharicum* Guerke.

*O. kilimandscharicum* Guerke is a valuable medicinal plant used both in traditional and modern medicine. The plant has neuro-protective [23], gastro-protective, digestive, antidiarrheal [24], carminative [25], antioxidant [26], antibacterial [26] antifungal [26], antipyretic [25], stimulant [25], anti-parasitic [27-29], wound-healing [30], and immunomodulatory [31] properties. The EO of *O. kilimandscharicum* is widely used in modern perfumery and the pharmaceutical industry. It is also used as a flavoring agent, in local application on sprains, in diarrheal medications, and in oral and dental preparations [25]. The EO has potent anti-oxidant [32, 33], anti-inflammatory [33], anti-cancer [33], and pesticidal qualities [34-36]. The variations in the chemical composition of EO of *O. kilimandscharicum* populations grown in different parts of the world have been widely investigated [37, 38]. The main components of the EO include camphor, 1, 8-cineol,  $\alpha$ -pinene, linalool, limonene, eugenol, methyl chavicol,  $\alpha$ - and  $\beta$ -bisabolene [39, 40]. The present study shows the Ultraviolet (UV) and Fourier Transform Infrared (FTIR) spectra of *O. kilimandscharicum* Guerke EO, which show very high amounts of eugenol.

## 2. Material and Methods


### 2.1. Plant Materials

The fresh leaves and flowers of *O. kilimandscharicum* were collected from the cultivated habitats in the medicinal and herbal garden of the Pharmacy Department of Lovely Professional University in February 2019. The fresh plant materials

were separated into two different parts. The first part was immediately processed for EO extraction, and the second part was labeled and stored in a refrigerator ( $-20^{\circ}\text{C}$ ) for future use.

## 2.2. Pharmacognostic Evaluation

In order to ensure the plant identity and prevent adulteration, the collected plant materials were subjected to pharmacognostic evaluation. *O. kilimandscharicum* is an evergreen, aromatic, perennial undershrub with simple ovate-oblong leaves, light purple or white-colored flowers, and ovoid-oblong black or brown-colored mucilaginous seeds. The EO is a light-yellow liquid with a strong odor of camphor (Bha12) (Figure 1). Histological studies of *O. kilimandscharicum* leaves were characterized by the presence of isobilateral lamina covered with a cuticle layer; glandular trichomes with multicellular heads containing essential oil (especially newly collected leaves); multicellular hair-covered trichomes; midrib arc-shaped vascular bundles consisting of xylem and phloem; and 3–4 layers of collenchymatous tissue present on the upper side of the vascular bundles (Figure 2).



**Figure 2.** Microscopic illustration of *O. kilimandscharicum* in 10X magnification A. leaf cross section, B. Hairy trichomes, and C. Glandular trichomes.


## 2.3. Essential oil Extraction

For the extraction of the EO, an accurately weighed amount of the plant material was subjected to hydrodistillation for 3 hours using a Clevenger-type apparatus, as previously described [26]. After completing the distillation process, the apparatus was allowed to cool and form clear yellow EO above the water inside the graded tube of the distillation apparatus. The condensed oil was collected and dried over anhydrous sodium sulfate and then stored in the dark at  $-20^{\circ}\text{C}$ .

## 2.4. UV Analysis of Essential Oil

### 2.4.1. Preparation of Eugenol Stock Solution

The stock solution of eugenol (1 mg/ml) was prepared by dissolving the accurately measured eugenol in methanol. The standard working solutions of eugenol were diluted from prepared stock solutions. The sample solutions were prepared by diluting the EO in an appropriate solvent (methanol) at a ratio of 1:1200 [41]. To quantify the concentration of eugenol in the EO of *O. kilimandscharicum*, the eugenol standard curve was plotted (Figure 3).



**Figure 3.** Eugenol standard curve.

### 2.4.2. UV Scanning of Essential Oil

The molecular absorption Ultraviolet- visible (UV-Vis) spectra of the EO of plant materials were recorded in the 200–400nm region. A computer-operated spectrophotometer was used with a 1 cm optical path quartz cuvette. The data were acquired and processed using the software package (UV Probe, Ver. 2.61). The same protocol was followed for the eugenol standards [42].


## 2.5. FTIR Analysis of Essential Oil

The FTIR spectra of the EO were acquired on a Shimadzu FTIR Spectrometer 8400S. The environmental conditions were maintained (25°C temperature and 30% humidity). The signal-to-noise ratio was 20000:1. Ten percent of the EO and reference chemicals and their specific solvents were deposited directly onto the Attenuated Total Reflection (ATR) sampling device without any treatment. The corresponding spectra were recorded in the range of 400–500  $\text{cm}^{-1}$  with a resolution of 4  $\text{cm}^{-1}$  and 12 scans for the sample and background. The velocity of the scan was 10 kHz with an interferogram size of 14,220 points. Before scanning each sample, the device was cleaned with 70% ethanol, and the background of air was taken. Shimadzu's IRsolution software was used to acquire spectra of samples and spectral manipulation [43, 44].


## 3. Result and Discussion

### 3.1. UV Analysis of Essential Oil

The UV spectra of standard eugenol and EO of *O. kilimandscharicum* are presented below (Figure 4 and Figure 5). In both spectra, eugenol has shown a peak absorbance at 282 nm. Based on the plotted standard curve of eugenol, the EO of *O. kilimandscharicum* contained 0.19  $\mu\text{g}/\text{ml}$  of eugenol.




**Figure 4.**  
The UV absorption spectrum of EO of *O. kilimandscharicum*.




**Figure 5.**  
The UV absorption spectrum of eugenol.

### 3.2. FTIR Analysis of Essential Oil

In this study, the FTIR absorption of EO, which was extracted from the leaves and flowers of *O. kilimandscharicum*, was investigated. The FTIR spectra of the EO and eugenol are shown below (Figure 6 and Figure 7). The presence of different functional groups was determined by comparing the vibration frequencies in wave numbers of the EO and eugenol spectra with those of an IR correlation chart (Table 1).



**Figure 6.**  
FTIR spectrum of EO of *O. kilimandscharicum*.



**Figure 7.**  
FTIR spectrum of Eugenol.

Based on the recorded FTIR spectra, eugenol showed its signature peaks at  $2968.55\text{ cm}^{-1}$ ,  $2841.24\text{ cm}^{-1}$ ,  $1435.09\text{ cm}^{-1}$ , and  $441.71\text{ cm}^{-1}$ , which were completely and/or partially different from the one of EO, but the other peaks were the same for both the eugenol and the EO. In the FTIR absorption spectra of EO and eugenol, the occurrence of peaks in the region  $3522.13\text{ cm}^{-1}$  and  $3516.35\text{ cm}^{-1}$  indicate the presence of hydroxyl groups, which may correspond to alcohols. The presence of peaks within the regions  $2968.55\text{ cm}^{-1}$  (C–H),  $2355.16\text{ cm}^{-1}$ ,  $2960.83\text{ cm}^{-1}$ ,  $2357.09\text{ cm}^{-1}$ ,  $2841.24\text{ cm}^{-1}$ ,  $1367.58\text{ cm}^{-1}$ ,  $1442.8\text{ cm}^{-1}$ ,  $1365.65\text{ cm}^{-1}$ ,  $1435.09\text{ cm}^{-1}$  (CH<sub>3</sub> group) plus a band at  $744.55\text{ cm}^{-1}$  (–CH<sub>2</sub> group), and  $644.25\text{ cm}^{-1}$ – $646.17\text{ cm}^{-1}$  is indicative of saturated aliphatic structures. The sharp peaks observed in the regions  $1606.76\text{ cm}^{-1}$  and  $1510.31\text{ cm}^{-1}$  may show the stretching of C=C of the aromatic moiety. The existence of halogenated aliphatic compounds may easily be determined from the peaks within the regions  $553.59\text{ cm}^{-1}$ ,  $594.1\text{ cm}^{-1}$ , and  $596.02\text{ cm}^{-1}$  [45].



Although the EO of most of the species of genus *Ocimum*, including *O. kilimandscharicum*, are valuable sources of eugenol, the main phytoconstituents for which they are known may vary [46, 47]. This phytochemical polymorphism even exists within the same plant populations [26, 38].

There could be several reasons for the profile polymorphism of *O. kilimandscharicum* EO viz, climate [48] or environmental factors [49-51], geographical zones [52-54], season [55-57], plant ontogeny [58, 59], harvesting time [60], plant chemo-type [26], stress factors [61, 62], genetic variability [26], etc.

**Table 1.**  
IR correction chart.

Frequencies on FTIR Spectra (1/cm)		Assignment of bonds	Mode of Vibration
<i>O. kilimandscharicum</i> oil	Eugenol		
3522.13	3516.35	O-H	Stretching
-	2968.55	C-H (hydrocarbons)	Stretching
2355.16, 2960.83	2357.09, 2841.24	CH <sub>3</sub>	Stretching
1606.76	1606.76	C=C	Stretching
1510.31	1510.31	C=C	Stretching
1367.58, 1442.8	1365.65, 1435.09	C-H (methyl)	Rocking
1265.35	1265.35	C-O/C-N	Stretching
1222.91	1224.84	C-N	Stretching
1030.95, 1136.11	1030.02, 1136.11	C-O/C-N	Stretching
995.3	997.23	=C-H	Bending
910.43	910.43	=C-H	Bending
804.34, 852.56	804.34, 852.56	C-H	Oop
744.55	744.55	-C-H (methylene)	Rocking
644.25	646.17	C-H	Bending
553.59, 594.1	553.59, 596.02	C-X	Stretching
435.93	441.71		

Several physicochemical techniques help assess and quantify the chemical fingerprint of EOs and their main phytoconstituents [63]. The UV and FTIR spectrometry techniques are two examples of tools currently used for these purposes [64, 65].

After identifying the main components of the EO, the UV spectrum could be used as a valuable tool for comparing and quantifying the desired phytoconstituents of the EO with the available standard. In this study, eugenol showed its peak absorbance around 282 nm in both the EO and pure eugenol spectra. Additionally, the EO contained 0.19 µg/ml of eugenol. Moreover, the availability of the FTIR fingerprint of every single chemotype would assist researchers in classifying the populations of a plant in specific clusters, based on the main phytoconstituents, and assess the purity of their EOs.

#### 4. Conclusion

*Ocimum kilimandscharicum* is an important medicinal plant of the genus *Ocimum*, which shows a high level of phytochemical polymorphism. The EO of different plant populations possesses various types of phytoconstituents of medicinal and economic value. Eugenol is one of these components present in high amounts in the EO of *O. kilimandscharicum*. The UV and FTIR spectra of the EO confirmed this as the eugenol showed peak absorption at 282 nm in the UV region and an intense vibration frequency in the FTIR spectrum obtained. Thus, the UV and FTIR fingerprint of EOs may consider as a rapid and reliable tool for the preliminary assessment and quantification of different phytoconstituents in the EOs.

#### References

- [1] T. Koeduka, E. Fridman, D. R. Gang, D. G. Vassão, B. L. Jackson, C. M. Kish, and J. P. Noel, "Eugenol and isoeugenol, characteristic aromatic constituents of spices, are biosynthesized via reduction of a coniferyl alcohol ester," *Proceedings of the National Academy of Sciences*, vol. 103, pp. 10128-10133, 2006. Available at: <https://doi.org/10.1073/pnas.0603732103>.
- [2] K. H. Tan and R. Nishida, "Methyl eugenol: Its occurrence, distribution, and role in nature, especially in relation to insect behavior and pollination," *Journal of Insect Science*, vol. 12, pp. 1-60, 2012. Available at: <https://doi.org/10.1673/031.012.5601>.
- [3] G. İlhami, "Antioxidant activity of eugenol: A structure-activity relationship study," *Journal of Medicinal Food*, vol. 14, pp. 975-985, 2011. Available at: <https://doi.org/10.1089/jmf.2010.0197>.
- [4] F. R. Giuliani, "The composition, structure, sources, and applications of Eugenol," *ESSAI*, vol. 12, pp. 1- 4, 2014.
- [5] B. Pavithra, "Eugenol-a review," *Journal of Pharmaceutical Sciences and Research*, vol. 6, pp. 153-154, 2014.
- [6] C.-K. Park, K. Kim, S. J. Jung, M. J. Kim, D. K. Ahn, S.-D. Hong, and S. B. Oh, "Molecular mechanism for local anesthetic action of eugenol in the rat trigeminal system," *Pain®*, vol. 144, pp. 84-94, 2009. Available at: <https://doi.org/10.1016/j.pain.2009.03.016>.

- [7] F. F. M. Da Silva, F. J. Q. Monte, T. L. G. de Lemos, P. G. G. Do Nascimento, A. K. de Medeiros Costa, and L. M. M. De Paiva, "Eugenol derivatives: Synthesis, characterization, and evaluation of antibacterial and antioxidant activities," *Chemistry Central Journal*, vol. 12, pp. 1-9, 2018. Available at: <https://doi.org/10.1186/s13065-018-0407-4>.
- [8] J. N. Barboza, C. D. S. M. Bezerra Filho, R. O. Silva, J. V. R. Medeiros, and D. P. de Sousa, "An overview on the anti-inflammatory potential and antioxidant profile of Eugenol," *Oxidative Medicine and Cellular Longevity*, vol. 2018, p. 3957262, 2018. Available at: <https://doi.org/10.1155/2018/3957262>.
- [9] F. Benencia and M. Courreges, "In vitro and in vivo activity of eugenol on human herpesvirus," *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, vol. 14, pp. 495-500, 2000. Available at: [https://doi.org/10.1002/1099-1573\(200011\)14:7%3C495::aid-ptr650%3E3.0.co;2-8](https://doi.org/10.1002/1099-1573(200011)14:7%3C495::aid-ptr650%3E3.0.co;2-8).
- [10] S. M. Ali, A. A. Khan, I. Ahmed, M. Musaddiq, K. S. Ahmed, H. Polasa, and N. Ahmed, "Antimicrobial activities of Eugenol and Cinnamaldehyde against the human gastric pathogen *Helicobacter pylori*," *Annals of Clinical Microbiology and Antimicrobials*, vol. 4, pp. 1-7, 2005. Available at: <https://doi.org/10.1186/1476-0711-4-20>.
- [11] R. M. Abd El-Baky and Z. S. Hashem, "Eugenol and linalool: Comparison of their antibacterial and antifungal activities," *African Journal of Microbiology Research*, vol. 10, pp. 1860-1872, 2016. Available at: <https://doi.org/10.5897/ajmr2016.8283>.
- [12] M. Asha, D. Prashanth, B. Murali, R. Padmaja, and A. Amit, "Anthelmintic activity of essential oil of *Ocimum sanctum* and eugenol," *Fitoterapia*, vol. 72, pp. 669-670, 2001. Available at: [https://doi.org/10.1016/s0367-326x\(01\)00270-2](https://doi.org/10.1016/s0367-326x(01)00270-2).
- [13] A. M. El-Kady, A. A. Ahmad, T. M. Hassan, H. E. El-Deek, S. S. Fouad, and S. S. Althagfan, "Eugenol, a potential schistosomicidal agent with anti-inflammatory and antifibrotic effects against *Schistosoma mansoni*, induced liver pathology," *Infection and Drug Resistance*, vol. 12, pp. 709-719, 2019. Available at: <https://doi.org/10.2147/idr.s196544>.
- [14] G. Chung and S. B. Oh, "Eugenol as local anesthetic. In K. Ramawat, & J. M. Merillon (Eds.), *Natural Products*," ed Heidelberg, Berlin: Springer, 2013, pp. 4001- 4015.
- [15] E. Omer, H. Said-Al Ahl, and S. Hendawy, "Production, chemical composition and volatile oil of different basil species/varieties cultivated under Egyptian soil salinity conditions," *Research Journal of Agriculture and Biological Sciences*, vol. 4, pp. 293-300, 2008.
- [16] A. K. Anand, M. Mohan, S. Z. Haider, and A. Sharma, "Essential oil composition and antimicrobial activity of three *Ocimum* species from Uttarakhand (India)," *International Journal of Pharmacy and Pharmaceutical Sciences*, vol. 3, pp. 223-225, 2011.
- [17] C. Kashyap, K. Ranjeet, A. Vikrant, and K. Vipin, "Therapeutic potency of *Ocimum kilimandscharicum* Guerke-a review," *Global Journal of Pharmacology*, vol. 5, pp. 191-200, 2011.
- [18] A. Viña and E. Murillo, "Essential oil composition from twelve varieties of basil (*Ocimum* spp) grown in Colombia," *Journal of the Brazilian Chemical Society*, vol. 14, pp. 744-749, 2003. Available at: <https://doi.org/10.1590/s0103-50532003000500008>.
- [19] K. Carović-Stanko, S. Orlić, O. Politeo, F. Strikić, I. Kolak, M. Milos, and Z. Satovic, "Composition and antibacterial activities of essential oils of seven *Ocimum* taxa," *Food Chemistry*, vol. 119, pp. 196-201, 2010. Available at: <https://doi.org/10.1016/j.foodchem.2009.06.010>.
- [20] R. S. Verma, R. C. Padalia, A. Chauhan, and S. T. Thul, "Exploring compositional diversity in the essential oils of 34 *Ocimum* taxa from Indian flora," *Industrial Crops and Products*, vol. 45, pp. 7-19, 2013. Available at: <https://doi.org/10.1016/j.indcrop.2012.12.005>.
- [21] A. S. Costa, M. D. Arrigoni-Blank, J. L. Carvalho Filho, A. D. Santana, D. D. Santos, P. B. Alves, and A. F. Blank, "Chemical diversity in Basil (*Ocimum* sp) Germplasm," *The Scientific World Journal*, vol. 2015, p. 352638, 2015. Available at: <https://doi.org/10.1155/2015/352638>.
- [22] D. J. Charles and J. E. Simon, "Essential oil constituents of *Ocimum kilimandscharicum* Guerke," *Journal of Essential Oil Research*, vol. 4, pp. 125-128, 1992.
- [23] V. Singh, P. Krishan, N. Singh, A. Kumar, and R. Shri, "Amelioration of ischemia-reperfusion induced functional and biochemical deficit in mice by *Ocimum kilimandscharicum* leaf extract," *Biomedicine & Pharmacotherapy*, vol. 85, pp. 556-563, 2017. Available at: <https://doi.org/10.1016/j.biopha.2016.11.064>.
- [24] R. V. Sarin, S. Narwal, and P. A. Bafna, "Anti-diarrhoeal activity of aqueous extract of *Ocimum kilimandscharicum*," *Journal of Ethnopharmacology*, vol. 148, pp. 223-228, 2013. Available at: <https://doi.org/10.1016/j.jep.2013.03.083>.
- [25] M. Bhasin, "Ocimum-taxonomy, medicinal potentialities and economic value of essential oil," *Journal of Biosphere*, vol. 1, pp. 48-50, 2012.
- [26] T. Chaturvedi, A. Kumar, A. Kumar, R. S. Verma, R. C. Padalia, V. Sundaresan, and K. Venkatesha, "Chemical composition, genetic diversity, antibacterial, antifungal and antioxidant activities of camphor-basil (*Ocimum kilimandscharicum* Guerke)," *Industrial Crops and Products*, vol. 118, pp. 246-258, 2018. Available at: <https://doi.org/10.1016/j.indcrop.2018.03.050>.
- [27] E. J. Kweka, E. E. Lyatuu, M. A. Mboya, B. J. Mwang'onde, and A. M. Mahande, "Oviposition deterrence induced by *Ocimum kilimandscharicum* and *Ocimum suave* extracts to gravid *Anopheles gambiae* ss (Diptera: Culicidae) in laboratory," *Journal of Global Infectious Diseases*, vol. 2, pp. 242-245, 2010. Available at: <https://doi.org/10.4103/0974-777x.68524>.
- [28] B. Jembere, D. Obeng-Ofori, A. Hassanali, and G. Nyamasyo, "Products derived from the leaves of *Ocimum kilimandscharicum* (Labiatae) as post-harvest grain protectants against the infestation of three major stored product insect pests," *Bulletin of Entomological Research*, vol. 85, pp. 361-367, 1995. Available at: <https://doi.org/10.1017/s0007485300036099>.

- [29] E. J. Kweka, H. M. Nkya, L. Lyaruu, E. E. Kimaro, B. J. Mawang'onde, and A. M. Mahande, "Efficacy of *Ocimum kilimandscharicum* plant extracts after four years of storage against *Anopheles gambiae* ss," *Journal of Cell and Animal Biology*, vol. 3, pp. 171-174, 2009.
- [30] P. S. Mahesh, M. Patil, R. Kumar, and S. R. Patil, "Evaluation of aqueous extract of leaves of *Ocimum kilimandscharicum* on wound healing activity in albino wistar rats," *International Journal of Pharmtech Research*, vol. 1, pp. 544-550, 2009.
- [31] G. Nahak and R. Sahu, "Immunomodulatory activity of aqueous leaf extract of *Ocimum kilimandscharicum* Guerke in *Clarias batrachus*," *International Journal of Pharmaceutical Sciences Review and Research*, vol. 28, pp. 137-146, 2014.
- [32] S. K. Singh, A. Anand, S. K. Verma, M. A. Siddiqui, A. Mathur, and S. Saklan, "Analysis of phytochemical and antioxidant potential of *Ocimum kilimandscharicum* Linn," *International Current Pharmaceutical Research Journal*, vol. 3, pp. 40-46, 2011.
- [33] V. De Lima, M. Vieira, C. Kassuya, C. Cardoso, J. Alves, M. Foglio, and A. Formagio, "Chemical composition and free radical-scavenging, anticancer and anti-inflammatory activities of the essential oil from *Ocimum kilimandscharicum*," *Phytomedicine*, vol. 21, pp. 1298-1302, 2014. Available at: <https://doi.org/10.1016/j.phymed.2014.07.004>.
- [34] D. Obeng-Ofori, C. Reichmuth, A. Bekele, and A. Hassanali, "Toxicity and protectant potential of camphor, a major component of essential oil of *Ocimum kilimandscharicum*, against four stored product beetles," *International Journal of Pest Management*, vol. 44, pp. 203-209, 1998. Available at: <https://doi.org/10.1080/096708798228112>.
- [35] J. Bekele and A. Hassanali, "Blend effects in the toxicity of the essential oil constituents of *Ocimum kilimandscharicum* and *Ocimum kenyense* (Labiateae) on two post-harvest insect pests," *Phytochemistry*, vol. 57, pp. 385-391, 2001. Available at: [https://doi.org/10.1016/s0031-9422\(01\)00067-x](https://doi.org/10.1016/s0031-9422(01)00067-x).
- [36] K. Chowdhary, A. Kumar, S. Sharma, R. Pathak, and M. Jangir, "*Ocimum* sp.: Source of biorational pesticides," *Industrial Crops and Products*, vol. 122, pp. 686-701, 2018. Available at: <https://doi.org/10.1016/j.indcrop.2018.05.068>.
- [37] V. Singh, P. Krishan, and R. Shri, "*Ocimum kilimandscharicum* Guerke: Phytochemical and pharmacological aspects; A review," *Journal of Pharmacognosy and Phytochemistry*, vol. 3, pp. 1-11, 2014.
- [38] L. Ntezurbanza, J. Scheffer, A. Looman, and A. B. Svendsen, "Composition of essential oil of *Ocimum kilimandscharicum* Grown in Rwanda1," *Planta Medica*, vol. 50, pp. 385-388, 1984. Available at: <https://doi.org/10.1055/s-2007-969744>.
- [39] R. Joshi, "Chemical composition of the essential oil of camphor basil (*Ocimum kilimandscharicum* Guerke)," *Global Journal of Medicinal Plant Research*, vol. 1, pp. 207-209, 2013.
- [40] M. . Ampah, M. A. . Brago, M. A. . Boatemaa, P. . Arthur, and M. M. . Sakyiamah, "Extraction and Characterization of Chitin and Chitosan Biopolymer as Wound Healing Material Using Shrimp Shells", 57, vol. 10, no. 2, pp. 69-75, 2021. <https://doi.org/10.18488/journal.57.2021.102.69.75>
- [41] F. Inam, S. Deo, and N. Narkhede, "HPLC-UV method development and quantification of eugenol from methanolic extracts of some spices," *International Journal of Chemical and Physical Sciences*, vol. 6, pp. 96-102, 2014.
- [42] D. Kruzlikova, J. Mocak, E. Katsoyannos, and E. Lankmayr, "Classification and characterization of olive oils by UV-Vis absorption spectrometry and sensorial analysis," *Journal of Food and Nutrition Research*, vol. 47, pp. 181-188, 2008.
- [43] E. S. Madivoli, L. Gitu, and E. Gumba, "Isolation and identification of essential oils from *Cymbopogon citratus* (stapf) Dc Using GC-MS and FT-IR," *Chemistry and Materials Research*, vol. 2, pp. 13-23, 2012.
- [44] R. N. Oliveira, M. C. Mancini, F. C. S. D. Oliveira, T. M. Passos, B. Quilty, R. M. D. S. M. Thiré, and G. B. McGuinness, "FTIR analysis and quantification of phenols and flavonoids of five commercially available plants extracts used in wound healing," *Subject Rio De Janeiro*, vol. 21, pp. 767-779, 2016. Available at: <https://doi.org/10.1590/s1517-707620160003.0072>.
- [45] K. Pramod, C. V. Suneesh, S. Shanavas, S. H. Ansari, and J. Ali, "Unveiling the compatibility of eugenol with formulation excipients by systematic drug-excipient compatibility studies," *Journal of Analytical Science and Technology*, vol. 6, pp. 1-14, 2015. Available at: <https://doi.org/10.1186/s40543-015-0073-2>.
- [46] K. Bhuvaneshwari, A. Gokulanathan, M. Jayanthi, V. Govindasamy, L. Milella, S. Lee, and S. Girija, "Can *Ocimum basilicum* L. and *Ocimum tenuiflorum* L. in vitro culture be a potential source of secondary metabolites?," *Food Chemistry*, vol. 194, pp. 55-60, 2016. Available at: <https://doi.org/10.1016/j.foodchem.2015.07.136>.
- [47] D. R. Gang, J. Wang, N. Dudareva, K. H. Nam, J. E. Simon, E. Lewinsohn, and E. Pichersky, "An investigation of the storage and biosynthesis of phenylpropenes in sweet basil," *Plant Physiology*, vol. 125, pp. 539-555, 2001. Available at: <https://doi.org/10.1104/pp.125.2.539>.
- [48] M. Shams, S. Z. Esfahan, E. Z. Esfahan, H. N. Dashtaki, A. Dursun, and E. Yildirim, "Effects of climatic factors on the quantity of essential oil and dry matter yield of coriander (*Coriandrum sativum* L.)," *Indian Journal of Science and Technology*, vol. 9, pp. 1-4, 2016. Available at: <https://doi.org/10.17485/ijst/2016/v9i6/61301>.
- [49] M. Shohayeb, A. Hassan, E. Abdel-Hameed, and B. Saleh, "Effects of macro-and microelements in soil of rose farms in Taif on essential oil production by *Rosa damascena* Mill," *Journal of Chemistry*, vol. 2015, pp. 1- 7, 2015. Available at: <https://doi.org/10.1155/2015/935235>.
- [50] Y. Dizajeyekan, A. Haghighi, and T. Gajoti, "Regional altitude and soil physicochemical factors influence the essential oil of *Thymus pubescens* (Lamiaceae: Lamiaceae)," *Journal of Biological & Environmental Sciences*, vol. 10, pp. 45-51, 2016.
- [51] S. F. Hendawy, M. S. Hussein, H. M. Amer, A. E. El-Gohary, and W. S. Soliman, "Effect of soil type on growth, productivity, and essential oil constituents of rosemary, *Rosmarinus officinalis*," *Asian Journal of Agriculture and Biology*, vol. 5, pp. 303-311, 2017.



- [52] A. Sanli and T. Karadoğan, “Geographical impact on essential oil composition of endemic *Kundmannia anatolica* Hub.-Mor.(Apiaceae),” *African Journal of Traditional, Complementary and Alternative Medicines*, vol. 14, pp. 131-137, 2017. Available at: <https://doi.org/10.21010/ajtcam.v14i1.14>.
- [53] L. C. Soro, S. Munier, Y. Pelissier, L. Grosmaire, R. Yada, D. Kitts, and C. Menut, “Influence of geography, seasons and pedology on chemical composition and anti-inflammatory activities of essential oils from *Lippia multiflora* Mold leaves,” *Journal of Ethnopharmacology*, vol. 194, pp. 587-594, 2016. Available at: <https://doi.org/10.1016/j.jep.2016.10.047>.
- [54] I. Almas, E. Innocent, F. Machumi, and W. Kisinza, “Effect of geographical location on yield and chemical composition of essential oils from three *Eucalyptus* species growing in Tanzania,” *Traditional Asian Medicine*, vol. 14, pp. 1-12, 2018.
- [55] G. Kofidis, A. Bosabalidis, and S. Kokkini, “Seasonal variation of essential oils in a linalool-rich chemotype of *Mentha spicata* grown wild in Greece,” *Journal of Essential Oil Research*, vol. 16, pp. 469-472, 2004. Available at: <https://doi.org/10.1080/10412905.2004.9698773>.
- [56] D. P. Costa, S. C. Santos, J. C. Seraphin, and P. H. Ferri, “Seasonal variability of essential oils of *Eugenia uniflora* leaves,” *Journal of the Brazilian Chemical Society*, vol. 20, pp. 1287-1293, 2009. Available at: <https://doi.org/10.1590/s0103-50532009000700013>.
- [57] S. D. Sá, T. S. Fiuza, L. L. Borges, H. D. Ferreira, L. M. Tresvenzol, P. H. Ferri, and J. R. Paula, “Chemical composition and seasonal variability of the essential oils of leaves and morphological analysis of *Hyptis carpinifolia*,” *Brazilian Journal of Pharmacognosy*, vol. 26, pp. 688-693, 2016. Available at: <https://doi.org/10.1016/j.bjp.2016.05.011>.
- [58] R. Croteau, M. Felton, F. Karp, and R. Kjonaas, “Relationship of camphor biosynthesis to leaf development in sage (*Salvia officinalis*),” *Plant Physiology*, vol. 67, pp. 820-824, 1981. Available at: <https://doi.org/10.1104/pp.67.4.820>.
- [59] R. P. Rocha, E. De Castro Melo, L. C. Barbosa, A. Almeida, R. H. S. Dos Santos, P. R. Cecon, and A. Santi, “Influence of plant age on the content and composition of essential oil of *Cymbopogon citratus* (DC.) Stapf,” *Journal of Medicinal Plants Research*, vol. 8, pp. 1121-1126, 2014. Available at: <https://doi.org/10.5897/jmpr2013.5549>.
- [60] V. D. Zheljzkov, C. L. Cantrell, T. Astatkie, and M. W. Ebelhar, “Peppermint productivity and oil composition as a function of nitrogen, growth stage, and harvest time,” *Agronomy Journal*, vol. 102, pp. 124-128, 2010. Available at: <https://doi.org/10.2134/agronj2009.0256>.
- [61] D. R. Gouvea, L. Gobbo-Neto, and N. P. Lopes, “The influence of biotic and abiotic factors on the production of secondary metabolites in medicinal plants,” *Plant Bioactives and Drug Discovery: Principles, Practice, and Perspectives*, vol. 17, pp. 419-452, 2012. Available at: <https://doi.org/10.1002/9781118260005.ch12>.
- [62] N. A. Abdelmajeed, E. N. Danial, and H. S. Ayad, “The effect of environmental stress on qualitative and quantitative essential oil of aromatic and medicinal plants,” *Archives Des Sciences*, vol. 66, pp. 100-120, 2013.
- [63] L.-H. Wang and W.-C. Sung, “Rapid evaluation and quantitative analysis of eugenol derivatives in essential oils and cosmetic formulations on human skin using attenuated total reflectance–infrared spectroscopy,” *Journal of Spectroscopy*, vol. 26, pp. 43–52, 2011. Available at: <https://doi.org/10.3233/SPE-2011-0526>.
- [64] P. Hernández-Sánchez, S. López-Miranda, C. Lucas-Abellán, and E. Núñez-Delicado, “Complexation of Eugenol (EG), as main component of clove oil and as pure compound, with  $\beta$ - and HP- $\beta$ -CDs,” *Food and Nutrition Sciences*, vol. 3, pp. 716-723, 2012. Available at: <https://doi.org/10.4236/fns.2012.36097>.
- [65] A. Yugatama, S. Rohmani, and R. Apriliani, “Analysis of Eugenol content in ethanolic extract of galangal rhizome (*Alpinia galanga* L. Willd) ointment using UV-VIS spectrophotometry method,” presented at the 1st Muhammadiyah International Conference on Health and Pharmaceutical Development (MICH-PhD 2018), (pp. 90-95). SCITEPRESS- Science and Technology Publications, Lda, 2021.